

erably be taken at least 30 minutes before meals. Peak concentrations in plasma are attained in about 1 to 2 hours and after a 500-mg oral dose are reported to range from 3 to 6 micrograms/mL.

Peak plasma concentrations of ampicillin after a 500-mg intramuscular dose given as the sodium salt occur within about 1 hour and are reported to range from 7 to 14 micrograms/mL.

Ampicillin is widely distributed and therapeutic concentrations can be achieved in ascitic, pleural, and joint fluids. It crosses the placenta and small amounts are distributed into breast milk. There is little diffusion into the CSF except when the meninges are inflamed. About 20% is bound to plasma proteins and the plasma half-life is about 1 to 1.5 hours, but this may be increased in neonates, the elderly, and patients with renal impairment; in severe renal impairment half-lives of 7 to 20 hours have been reported.

Ampicillin is metabolised to some extent to penicilloic acid which is excreted in the urine.

Renal clearance of ampicillin occurs partly by glomerular filtration and partly by tubular secretion; it is reduced by probenecid. About 20 to 40% of an oral dose may be excreted unchanged in the urine in 6 hours; urinary concentrations have ranged from 0.25 to 1 mg/mL after a dose of 500 mg. After parenteral use about 60 to 80% is excreted in the urine within 6 hours. Ampicillin is removed by haemodialysis. High concentrations are reached in bile; it undergoes enterohepatic recycling and some is excreted in the faeces.

Ampicillin with sulbactam. The pharmacokinetics of ampicillin and sulbactam are broadly similar and neither appears to affect the other to any great extent.

Uses and Administration

Ampicillin is used in the treatment of a variety of infections due to susceptible organisms (see Antimicrobial Action, above). They include biliary-tract infections, bronchitis, endocarditis, gastro-enteritis (including salmonella enteritis and shigellosis), gonorrhoea, listeriosis, meningitis, perinatal streptococcal infections (intrapartum prophylaxis against group B streptococci), peritonitis, pneumonia, septicaemia, typhoid and paratyphoid fever, and urinary-tract infections. Resistance to ampicillin is increasingly a problem in some infections, for example, gonorrhoea, pneumococcal infections, respiratory-tract infections due to *Haemophilus influenzae* or *Moraxella catarrhalis* (*Branhamella catarrhalis*), *Salmonella* infections, shigellosis, and infections due to *Escherichia coli*. For details of these infections and their treatment, see under Choice of Antibacterial, p.162. If beta-lactamase-producing organisms are present, ampicillin can be given with a beta-lactamase inhibitor such as sulbactam (see below) or a penicillinase-resistant drug such as cloxacillin, dicloxacillin, or flucloxacillin (known as co-fluampicil). It may also be used with an aminoglycoside to increase the spectrum of organisms covered; it is advisable to give the injections separately.

Administration and dosage. The dosage of ampicillin will depend on the severity of the disease, the age of the patient, and renal function. Ampicillin is usually given orally as the trihydrate and by injection as the sodium salt. Doses are expressed in terms of the equivalent amount of ampicillin; 1.06 g of ampicillin sodium and 1.15 g of ampicillin trihydrate are each equivalent to about 1 g of ampicillin.

The usual adult oral dose is 0.25 to 1 g every 6 hours taken at least 30 minutes before or 2 hours after food. Children may be given half the adult dose. The usual adult dose by injection is 500 mg every 4 to 6 hours intramuscularly or by slow intravenous injection over

3 to 5 minutes or by infusion. Again, children may be given half the adult dose.

For urinary-tract infections, ampicillin 500 mg is given orally every 8 hours.

For typhoid and paratyphoid fever where *Salmonella typhi* strains remain sensitive to ampicillin, an oral dose of 1 to 2 g may be given every 6 hours for 2 weeks for acute infections, and for 4 to 12 weeks in carriers. An intramuscular dose of 10 mg/kg (maximum dose 250 mg) every 6 hours for 4 to 6 weeks has been suggested for children who are chronic carriers.

Ampicillin 2 g given with probenecid 1 g, as a single oral dose, has been used in the treatment of uncomplicated gonorrhoea in areas where gonococci remain sensitive; repeated doses are recommended in females.

In meningitis, higher parenteral doses of 2 to 3 g given intravenously every 4 or 6 hours have been suggested. For infants and children with meningitis, an intravenous dose of 150 mg/kg daily in divided doses may be given; a dose of 50 mg/kg (maximum 3 g) every 4 to 6 hours has also been suggested. Neonates may be given a dose of 50 mg/kg every 12 hours for those under 1 week of age, or every 8 hours for older neonates.

For intrapartum prophylaxis against group B streptococcal infection in the neonate, a maternal dose of 2 g by intravenous injection initially then 1 g every 4 hours until delivery has been suggested.

Ampicillin may also be given by other routes, usually as a supplement to systemic therapy. Intraperitoneal or intrapleural injections are given in a dose of 500 mg daily dissolved in 5 to 10 mL of water. For intra-articular injection, ampicillin 500 mg daily is given dissolved in up to 5 mL of water or a solution of procaine hydrochloride 0.5%.

Ampicillin benzathine has also been given by intramuscular injection.

Ampicillin with sulbactam. The sodium salts of ampicillin and sulbactam (p.335) may be given intramuscularly or intravenously in the treatment of infections due to beta-lactamase-producing organisms. Doses are expressed in terms of the equivalent amounts of ampicillin and sulbactam; available injections contain ampicillin and sulbactam in the ratio 2:1, respectively. The usual dose is ampicillin 1 g with sulbactam 500 mg every 6 hours; doses may be doubled in severe infections.

For oral use sultamicillin (p.344), a mutual prodrug of ampicillin and sulbactam, may be given.

Administration in renal impairment. The dose of ampicillin should be reduced, or the dose interval increased, in severe renal impairment (creatinine clearance less than 10 mL/minute). Patients undergoing dialysis should receive an additional dose after the session.

Preparations

BP 2008: Ampicillin Capsules; Ampicillin Injection; Ampicillin Oral Suspension; Co-fluampicil Capsules; Co-fluampicil Oral Suspension; **USP 31:** Ampicillin and Probenecid for Oral Suspension; Ampicillin and Sulbactam for Injection; Ampicillin Capsules; Ampicillin for Injectable Suspension; Ampicillin for Injection; Ampicillin for Oral Suspension; Ampicillin Tablets.

Proprietary Preparations (details are given in Part 3)

Arg.: Alpovex; Aminoxidin†; Ampi-Bis; Ampi†; Ampidler; Ampigen; Ampigrand; Ampinov; Ampitenk; Ampixen; Atecilina; Baclitina; Decilina†; Fabopicina; Galicidina†; Grampenil†; Histopen; Poenbiotico†; Trifacilina; Trimicro; Welticline. **Austral.:** Alphacin; Ampicyn; Astrapren; Ibimycin. **Austria:** Standacillin. **Belg.:** Pentrexyl. **Braz.:** Ampi; Ampicilan; Ampicikyna†; Ampicil; Ampicilab; Ampicilase; Ampicilbt; Ampicil†; Ampicilon; Ampicimax†; Ampicinal†; Ampifar†; Ampigran; Ampilozin; Ampitrat†; Ampival†; Ampivin†; Amplacina; Amplacin; Amplibac; Amplofen†; Bacterin†; Binotal; Bi-pencil†; Cililon; Clippen; Emicillin; Expectocilin; Gomot; Gramicilina; Lifacilina†; Notacilin†; Pratilicin†; Tandrexin†. **Canad.:** Apo-Amp; Nu-Amp. **Cz.:** Apo-Amp†; Penstabil†; Standacillin†. **Denn.:** Doktacillin†; Pentrexyl†. **Fin.:** A-Peni. **Fr.:** Totaper. **Ger.:** Binotal. **Gr.:** Copericlex; Isticiline; Pentrexyl. **Hong Kong:** Amprexyl; Dhacillin†; Pamecil; Penbritin†; Pendil; Pentrexyl. **Hung.:** Penstabil†; Semicillin; Standacillin. **India:** Ampilin; Ampipen; Ampisyn; Anglopex; Bestilina-A; Binotal; Bremecina; Brupen; Deamicilina; Dibacilina; Diferin; Expicin; Flamicina; Gramipen; F; Idfacilina; Lampicin; Marovilina; Meprizina; Mexapin; Mibiot; Omnipen; Pebit; Penbriter; Pentiver; Pentrexyl; Procilia; Prodifer; Promecilina; Rayepent†; Sinaplin; Tronex; Yapamicin; Zumorin. **Neth.:** Pentrexyl†. **Norw.:** Pentrexyl. **Philippines:** Aldribid; Ampicil; Ampines; Bactimed; Clovillin; Ecorin; Exicillin; Gramic; Obocil; Panacta; Penbritin; Pentrexyl; Picaplin; Polypen; Rotocin; Shinapan; Vatacl. **Port.:** Ampilar; Cilin†; Estreptobiotico; Hiperbiotico; Hibperiobiotico Retard. **S.Afr.:** Ampimax; Ampiren; Be-Ampid; Penbritin; Peterite; Tericillin; Ranamp; Spectracil. **Singapore:** Ampilin; Dhacillin; Pamecil†; Standacillin. **Spain:** Ampipus; Antibipen; Britapen; Gobemicina; Nuvafen; Nuvapen. **Swed.:** Doktaclillin. **Thail.:** Amcillin†; Amilar; Ampicyn; Ampilin; Ampilin†; Ampira; Amprexyl; Ampro†; Eracillin; Penbritin; Pencotrex; Pentrexyl; Siampicil; Sumapen; Vacillin; Vicillin. **Turk.:** Alfasilin; Ampisid; Ampisina; Neosilin; Penbisin; Silina. **UAE:** Julphapen. **UK:** Magnapen; Penbritin; Rimacillin. **USA:** Principen. **Venez.:** Alampen; Ampen†; Ämpenina; Ampiga†; Ampilan; Arcocillin; Fibrapen†; Intrapen†; Neoampicil†.

Amplital; Amplizer; Ibimicynt†; Pentrexyl; **Malaysia:** Ampilin; Biocil†; Pamecil; Setcollin†; Standacillin; **Mex.:** Acilmel; Alphapen; Alvedrin; Am-An; Ambidin; Ambiosol; Ampex; Ampi-Quin; Ampi-Tecno; Ampibit; Ampicid†; Ampidratt†; Ampigran; Ampilon†; Ampimec; Ampiset†; Amprexyn; Amsapen; Anglopex; Bestilina-A; Binotal; Bremecina; Brupen; Deamicilina; Dibacilina; Diferin; Expicin; Flamicina; Gramipen; F; Idfacilina; Lampicin; Marovilina; Meprizina; Mexapin; Mibiot; Omnipen; Pebit; Penbriter; Pentiver; Pentrexyl; Procilia; Prodifer; Promecilina; Rayepent†; Sinaplin; Tronex; Yapamicin; Zumorin. **Neth.:** Pentrexyl†. **Norw.:** Pentrexyl. **Philippines:** Aldribid; Ampicil; Ampines; Bactimed; Clovillin; Ecorin; Exicillin; Gramic; Obocil; Panacta; Penbritin; Pentrexyl; Picaplin; Polypen; Rotocin; Shinapan; Vatacl. **Port.:** Ampilar; Cilin†; Estreptobiotico; Hiperbiotico; Hibperiobiotico Retard. **S.Afr.:** Ampimax; Ampiren; Be-Ampid; Penbritin; Peterite; Tericillin; Ranamp; Spectracil. **Singapore:** Ampilin; Dhacillin; Pamecil†; Standacillin. **Spain:** Ampipus; Antibipen; Britapen; Gobemicina; Nuvafen; Nuvapen. **Swed.:** Doktaclillin. **Thail.:** Amcillin†; Amilar; Ampicyn; Ampilin; Ampilin†; Ampira; Amprexyl; Ampro†; Eracillin; Penbritin; Pencotrex; Pentrexyl; Siampicil; Sumapen; Vacillin; Vicillin. **Turk.:** Alfasilin; Ampisid; Ampisina; Neosilin; Penbisin; Silina. **UAE:** Julphapen. **UK:** Magnapen; Penbritin; Rimacillin. **USA:** Principen. **Venez.:** Alampen; Ampen†; Ämpenina; Ampiga†; Ampilan; Arcocillin; Fibrapen†; Intrapen†; Neoampicil†.

Multi-ingredient: **Arg.:** Aminoxidin Sulbactam; Ampi-Bis Plus; Ampigen SB; Ampibenzatin Bronqual; Aseptobron Ampicilina†; Cronopen Balsamico; Grampenil Bronqual†; Meticol; Pixin; Unasyn; Unasyn†; **Austria:** Unasn; **Braz.:** Ambezetal; Amplotal; Benzotol; Combactan; Duraper; Optacil; Parenzym; Ampicilina; Sulbacter†; Unasn; Urobioptic†; Uropielon; **Chile:** Unasn; **Cz.:** Ampicloxt; Unasn; **Fr.:** Unacid; **Ger.:** Begalin-P. **Hong Kong:** Ampicloxt; APT-Ampicloxa; Parnedox; Rosclox; Unasn; **Hung.:** Unasn; **India:** Adillo; Ampillox; AmpiloX-LB; Amplus; Ampoxin; Ampoxin-LB; Campicillin Plus; Campillo; Clax; Megadox; Megalo LB; Megapen; Sulbacin; **Ir.:** Ampicloxt; **Israel:** Unasn; **Ital.:** Ampi-plus†; Ampilum; Bethacil; Diampicil†; Loridin; Unasn; **Jpn.:** Sulperazon†; Unasn; **Malaysia:** Sulbacin; Unasn; **Mex.:** Ampiclo-D; Anglotex; Bisolvon A; Brucilina; Brupen Compuesto; Diamprex; Doxapen; Muclolin A; Panac; Panac K; Pentibrom; Penidix; Pentrexyl; Expert; Unasn; **Philippines:** Unasn; **Pol.:** Unasn; **Rus.:** Oxam (Оксам)†; Sultasin (Сультасин); Unasn (Унасин); **S.Afr.:** Ampicid; Ampiclo; Apen; Cloxam; Megamox; Rancilos†; **Singapore:** Unasn; **Spain:** Ampivel; Espectral†; Espectrosina†; Gobemicina Retard; Retarpen; Retarpen Balsamico†; Retarpen Mucolito†; Ultrapen; Unasn; **Thail.:** Ampidlox Sulam; Unasn; Vicillin-S; **Turk.:** Azosilin; Combicid; Duobaktam; Duocid; Nobecid; Sulbakt; Sulcid; Sutaisid; Sutaisid; **UK:** Magnapen; **USA:** Unasn; **Venez.:** Ampibactan; Ampitrent†; Fipexim; Sinf; Unasn.

Apramycin (BAN, USAN, rINN)


47657; Apramicine; Apramycine; Apramycinum; EL-857; EL-857/820; Nebramycin Factor 2. 4-O-[*(2R,3R,4aS,6R,7S,8aR)-3-*Amino-6-(4-amino-4-deoxy- α -D-glucopyranosyloxy)-8-hydroxy-7-methylaminoperhydropyran-*[3,2-b]pyran-2-yl*]-2-deoxystreptamine.

Апрамицин

$C_21H_{41}N_5O_{11}$ = 539.6.

CAS — 37321-09-8.

ATC Vet — QA07AA92; QJ01GB90; QJ51GB90.



Apramycin Sulfate (rINN)

Apramycin Sulphate (BAN); Apramycine, Sulfate d'; Apramycin Sulfas; Apramycinsulfat; Apramysiinsulfat; Sulfato de apramicina.

Апрамицина Сульфат

$C_21H_{41}N_5O_{11,2}/H_2SO_4$ = 784.8.

CAS — 41194-16-5.

Pharmacopoeias. In BP(Vet).

BP(Vet) 2008 (Apramycin Sulphate). The sulfate of an antibiotic produced by certain strains of *Streptomyces tenebrarius* or by other means. The potency is not less than 430 units per mg, calculated with reference to the anhydrous substance. A light brown hygroscopic powder or granular material. Freely soluble in water; practically insoluble in alcohol, in acetone, in ether, and in methyl alcohol.

Profile

Apramycin is an aminoglycoside antibiotic used as the sulfate in veterinary practice for the treatment of susceptible infections.