

Preparations

BP 2008: Benzylamine Cream; Benzylamine Mouthwash; Benzylamine Oromucosal Spray.


Proprietary Preparations (details are given in Part 3)

Arg.: Actifedrin; Benicfen; Ernex; Sandival Desleble; **Austral.:** Difflam; Difflam Anti-inflammatory Throat Spray; Difflam Solution; **Austria:** Tantum; Tantumar; **Cz.:** Benflogin; Benzitrat; Ciflogex; Erdamint; Flogin-Ped[†]; Flago Rossi; Flogolab; Flogoral; Forengoral; Neoflogin; **Canad.:** Sun-Benzil; Tantum; **Cz.:** Rosalgin; Tantum; **Denm.:** Andolex; **Fr.:** Opalgyn; **Ger.:** Tantum Rosat; Tantum Verde; **Gr.:** Tantum; **Hong Kong:** Dantum; Difflam; Verax; **Hung.:** Tantum Verde; **Indon.:** Tanflex; Tantum; **Ir.:** Difflam; **Israel:** Easy gel; **Ital.:** Alloben; Benzirin; Ginesal; Lagin[†]; Multum; Sanfor Collutorio; Tantum; Verax; Xentafid; **Malaysia:** Difflam Anti-inflammatory Lozenges; Difflam Solution; **Mex.:** Arthrofen; Lonol; Vantal; **Neth.:** Tantum; **NZ.:** Difflam; **Philippines:** Difflam; **Pol.:** Hascosept; Tantum; **Port.:** Flogoral; Momen; Rosalgin; Tantum; Tantum Rosa; Tantum Verde; **Rus.:** Tantum Rose (Тантум Роза); Tantum Verde (Тантум Верде); **S.Afr.:** Andolex; **Singapore:** Difflam; **Spain:** Agilon; Fujigum; Rosalgin; Tantum; Tantum Verde; **Swed.:** Andolex; **Switz.:** Bucco-Tantum; **Thail.:** Difflam; **Turk.:** Benzidan; Tanflex; Tantum; Ternex; **UK:** Difflam; **Venez.:** Azutan; Bevi Dam; Biozendi; Flodont; Ginacol; Tantum; Tantum Verde; Zydar.

Multi-ingredient: **Arg.:** Buxex; Dresan Biotic[†]; Dresant[†]; Ernex Duo; Epectocural; Pentadent[†]; **Austral.:** Difflam Anti-inflammatory Lozenges with Cough Suppressant; Difflam Lozenges; Difflam Mouth Gel; Difflam-C; Loginc Rapid Relief; **Braz.:** Angino-Rub; **Hong Kong:** Difflam Anti-inflammatory Lozenges; Difflam Mouth Gel; Difflam-C; Loginc Rapid Relief; **Hung.:** Tantum Rosa; **Ital.:** Algolinsal[†]; Gola Action; Linea F; Mediplus; **Malaysia:** Difflam Anti-inflammatory Lozenges (with Antibacterial); Difflam Anti-Inflammatory Lozenges (with cough suppressant); Difflam Mouth Gel; Difflam-C; **Mex.:** Lonol Sport; **NZ.:** Difflam Cough; Difflam Mouth Gel; Difflam-C; **Port.:** Benoral; Gartun; Tantum Verde; **S.Afr.:** Andolex-C; **Singapore:** Difflam Anti-inflammatory Anti-Bacterial Lozenges; Difflam Mouth Gel; Difflam-C; **Spain:** Bristacilina Dental; Dolosartol[†]; Etermol Antitusivo; Mentamida; Prosturo; Tantum; Vincisetil Otico; **Turk.:** Kroboren; **Venez.:** Amicets; Gencivil Compuesto; Solunovar Compuesto.

Benzyl Nicotinate

Bensylnikotinat; Bentsylinikotinaatti; Benzil Nikotinat; Benzyl-Nicotinas; Nicotinato de bencilo. Benzyl pyridine-3-carboxylate. $C_{13}H_{11}NO_2 = 213.2$. CAS — 94-44-0.



Pharmacopoeias. In **Ger.**

Profile

Benzyl nicotinate is used in topical preparations as a rubefacient.

Preparations

Proprietary Preparations (details are given in Part 3)

Ger.: Leukoma-Aktiv-Rheumabadt[†]; Pernionin Thermo Teilbad; Pernionin Thermo Vollbad; Pykaryl T; Rubriment; Rubriment-BN.

Multi-ingredient: **Arg.:** Butidionat[†]; Oxa Sport; Pergalen; **Austria:** Ambenat; Derivon; Expecta-Balsam; Igutur-antirheumatische; Igutur-Rheumafuß; Methoneurin; Mobilisin plus; Rhenex; Rubizone; Rubizone-Rheumagel; Rubriment; Thermo-Rheumon; Thrombophob; **Braz.:** Etrat[†]; Trombophob; **Chile:** Bayro-Therm[†]; **Cz.:** Dol-Rubriment[†]; Rheuma-Salbe[†]; Rubriment-N[†]; Thermo-Rheumon[†]; **Fin.:** Trombosol; **Fr.:** Lumbaligne; **Ger.:** ABC Wärme-Salbe[†]; Ambene N; Arthrodestal N[†]; Auroanal; Thermo; Camphopin; Capsamol N[†]; Caye Rheuma-Balsam; Con-Select[†]; DoloVisano Salbe[†]; Enasex-N[†]; Forapin E[†]; Hot Thermo; Lomazell forte N[†]; Lumbinon Thermo[†]; mikanil[†]; Ostochont[†]; Pelvichthol N; Phardol Rheuma[†]; Phardol Wärme-Balsam[†]; Phlogont-Thermal; Praecordia S[†]; Rheubalmal Thermo[†]; Rheuma-Salbe N; Rheuma-Salbe[†]; Rheumasalbe[†]; Rheumasan N[†]; Rheumatis[†]; Rubriment; Rubriment-N[†]; Tachyng Campher Herzsalbe; Thermo-Menthoneurin[†]; Thermo-Rheumon N[†]; Thermo-Rheumon[†]; Thermosenex; Togal Mobil-Gel[†]; Warme-Gel[†]; zuk thermo[†]; **Gr.:** Bayolin; Ehrlich; Strifasian[†]; Thermo-Roipion; **Hong Kong:** Salome-thy; **Hung.:** Bayolin[†]; Thermo-Rheumon[†]; **India:** Beparine; K5 Hair Tincture[†]; Thrombophob; **Indon.:** Stop X; Thrombophob; Zeropain; **Ital.:** Salongas; Sloan; **Mex.:** Bayre Temo[†]; **Pol.:** Lumbolin; Thermo-Rheumon; **Port.:** Adrinex[†]; Balsamo Analgesico; Mediglanan; **Rus.:** Capiscam (Капискам); Heparin Ointment (Гепариновая Мазь); **Switz.:** Assan thermo; Demotherm Pommade contre le rhumatisme[†]; Dolo Demotherm; Forapin[†]; Histalgane; Marament-N; Thermocutan[†]; **Turk.:** Thermo-Doline; Thermo-Rheumon; Thermoflex; **UK:** Salonair[†]; **Venez.:** Ehrlich Balsamo.

Beta-aminopropionitrile


Aminopropionitrile; β-Aminopropionitrile; β-Aminopropionitrile; BAPN. 3-Aminopropionitrile.

Бета-аминопропионитрил

$C_3H_6N_2 = 70.09$.

CAS — 151-18-8 (beta-aminopropionitrile); 1119-28-4 (beta-aminopropionitrile fumarate).

ATC Vet — QM01AX91.




Profile

Beta-aminopropionitrile, a lysyl oxidase inhibitor, is an anti-inflammatory used as the fumarate in veterinary medicine for the treatment of tendonitis.

Bicifadine Hydrochloride

(USAN, rINN)
Bicifadina, hidrocloruro de; Bicifadine, Hydrochloride de; Bicifadini Hydrochloridum; CL-220075. (±)-1-p-Tolyl-3-azabicyclo[3.1.0]hexane hydrochloride.
Бицифадина Гидрохлорид
 $C_{12}H_{15}N.HCl = 209.7$.
CAS — 71195-57-8 (bicifadine); 66504-75-4 (bicifadine hydrochloride).




Profile

Bicifadine hydrochloride is a novel analgesic under investigation for the treatment of painful conditions including postoperative dental pain.

Bornyl Salicylate

Borneol Salicylate; Salicilato de bornilo. 2-Hydroxybenzoic acid 1,7,7-trimethylbicyclo[2.2.1]hept-2-yl ester.

Борнилсалацилат
 $C_{17}H_{22}O_3 = 274.4$.
CAS — 560-88-3.



Profile

Bornyl salicylate is a salicylic acid derivative that has been used topically in rubefacient preparations similarly to methyl salicylate (p.85) for the relief of pain in musculoskeletal and joint disorders.

Preparations


Proprietary Preparations (details are given in Part 3)

Multi-ingredient: **Ger.:** Forapin E[†]; **Switz.:** Forapin[†]; Hygiderm[†].

Bromfenac Sodium

(USAN, rINN)
AHR-10282; AHR-10282B; Bromfenac Sodium; Bromfenaco sódico; Natrii Bromfenacum. Sodium [2-amino-3-(*p*-bromobenzoyl)phenyl]acetate sesquihydrate.

Натрий Бромфенак
 $C_{15}H_{11}BrNNaO_3 \cdot 1/2H_2O = 383.2$.
CAS — 91714-94-2 (bromfenac); 91714-93-1 (bromfenac sodium); 120638-55-3 (bromfenac sodium).



(bromfenac)

Profile

Bromfenac sodium, a phenylacetic acid derivative related to diclofenac (p.44), is an NSAID (p.96). It is instilled twice daily as 0.1% eye drops for ocular pain and inflammation including post-operative inflammation in patients who have undergone cataract extraction. When used postoperatively, it may be started 24 hours after surgery and continued for the next 14 days.

It was formerly given orally in the management of acute pain but was withdrawn from the market after reports of severe and sometimes fatal hepatic failure.

Effects on the eyes. Severe corneal melting (ulceration) was seen in 3 patients after topical use of bromfenac sodium.¹ All patients recovered after bromfenac was withdrawn. Similar ef-

fects have been reported with other ophthalmic NSAID preparations; for further details see under Adverse Effects of Diclofenac, p.45.

1. Asai T, et al. Three cases of corneal melting after instillation of a new nonsteroidal anti-inflammatory drug. *Cornea* 2006; **25:** 224-7.

Preparations

Proprietary Preparations (details are given in Part 3)

Jpn: Bronnick; **USA:** Xibrom.

Bufexamac


(BAN, rINN)
Bufeksamaakkis; Bufexamaco; Bufexamacum; Bufexamák; Bufexamak. 2-(4-Butoxyphenyl)acetohydroxamic acid.

Буфексамак

$C_{12}H_{17}NO_3 = 223.3$.
CAS — 2438-72-4.

ATC — M01AB17; M02AA09.

ATC Vet — QM01AB17; QM02AA09.



Pharmacopoeias. In **Eur.** (see p.vii) and **Jpn.**

Ph. Eur. 6.2 (Bufexamac). A white or almost white, crystalline powder. Practically insoluble in water; soluble in dimethylformamide; slightly soluble in ethyl acetate and in methyl alcohol. Protect from light.

Profile

Bufexamac is an NSAID (p.96) that is applied topically in concentrations of 5% in various skin disorders. Stinging and burning may occur after application; hypersensitivity reactions have been reported.

Preparations

Proprietary Preparations (details are given in Part 3)

Arg.: Parfenac[†]; **Austral.:** Paraderm Plus; **Fr.:** Bufex; Bufexan; Droxaryl; Parfenac; **Belg.:** Bufexinet[†]; Droxaryl[†]; **Cz.:** Droxaryl[†]; **Fr.:** Parfenac; **Ger.:** Bufederm[†]; duraderm; Haemo-Elixir; Bufexamac; Jomax; Malipuran; Parfenac; Windol; **Ital.:** Fansmac; Viafen; **Neth.:** Droxaryl; Parfenac; **Port.:** Parfenac; **Switz.:** Parfenac.

Multi-ingredient: **Austral.:** Paraderm Plus; Resolve; **Austria:** Droxaryl; **Cz.:** Mastu S; **Ger.:** Faktu akut; Hamo-riatopharm N; Hamoagil plus; Mastu S; **Hong Kong:** Fungo Soothing Balm; Mastu S; **Hung.:** Mastu S; **NZ.:** Paraderm Plus; **Rus.:** Proctosan (Проктозан); **Thail.:** Mastu S.

Bumadizone Calcium

Bumadizone cálcica; Bumadizone Calcique; Calcii Bumadizonom. Calcium 2-(1,2-diphenylhydrazinocarbonyl)hexanoate hemihydrate.


Кальций Бумадизон

$(C_{19}H_{12}N_2O_3)_2Ca \cdot H_2O = 699.8$.

CAS — 3583-64-0 (bumadizone); 34461-73-9 (bumadizone calcium).

ATC — M01AB07.

ATC Vet — QM01AB07.



(bumadizone)

Profile

Bumadizone calcium is an NSAID (p.96) that is metabolised to phenylbutazone (p.117) and oxyphenbutazone (p.107). Use has been limited by the risk of agranulocytosis and other haematological adverse effects.

Preparations

Proprietary Preparations (details are given in Part 3)

Mex.: Desflam.