

Hydrocortisone Sodium Succinate

(BANM, INNM) \otimes

Cortisol Sodium Succinate; Hydrocortisone, Succinate Sodium d'; Hydrocortisoni Natrii Succinatis; Hydrokortyzonu bursztynianu sól sodowa; Succinato sódico de hidrocortisona. Hydrocortisone 21-(sodium succinate).

Гидрокортизона Натрия Сукцинат

 $C_{25}H_{33}NaO_8 = 484.5$.

CAS — 125-04-2.

ATC — A01AC03; A07EA02; C05AA01; D07AA02; H02AB09; S01BA02; S02BA01.

ATC Vet — Q01AC03; Q07EA02; QC05AA01; QD07AA02; QH02AB09; QS01BA02; QS02BA01.

Pharmacopoeias. In *Chin., Int., It., Jpn., Pol., and US.*

USP 31 (Hydrocortisone Sodium Succinate). A white or nearly white, odourless, hygroscopic, amorphous solid. Very soluble in water and in alcohol; very slightly soluble in acetone; insoluble in chloroform. Store in airtight containers. Protect from light.

Hydrocortisone Valerate (BANM, USAN, INNM) \otimes

Cortisol Valerate; Hydrocortisone, Valérat d'; Hydrocortison Valeras; Valerato de hidrocortisona. Hydrocortisone 17-valerate.

Гидрокортизона Валерат

 $C_{26}H_{38}O_6 = 446.6$.

CAS — 57524-89-7.

ATC — A01AC03; A07EA02; C05AA01; D07AA02; H02AB09; S01BA02; S02BA01.

ATC Vet — Q01AC03; Q07EA02; QC05AA01; QD07AA02; QH02AB09; QS01BA02; QS02BA01.

Pharmacopoeias. In *US.*

Adverse Effects, Treatment, Withdrawal, and Precautions

As for corticosteroids in general (see p.1490).

When applied topically, particularly to large areas, when the skin is broken, or under occlusive dressings, corticosteroids may be absorbed in sufficient amounts to cause systemic effects. Prolonged use of ophthalmic preparations containing corticosteroids has caused raised intra-ocular pressure and reduced visual function.

Effects on fluid and electrolyte balance. A report of marked hypokalaemia and hypomagnesaemia associated with high-dose intravenous hydrocortisone therapy in an alcoholic patient with suspected immune thrombocytopenia.¹ Cardiac arrhythmias developed, and prolonged infusion of magnesium and potassium was required to restore normal plasma concentrations.

1. Ramsahoye BH, et al. The mineralocorticoid effects of high dose hydrocortisone. *BMJ* 1995; **310**: 656-7.

Effects on the nervous system. For reports and comments on paraesthesia or perineal irritation associated with hydrocortisone sodium phosphate given intravenously, see p.1492.

Hypersensitivity and anaphylaxis. References to hypersensitivity reactions and anaphylaxis associated with the intravenous use of hydrocortisone;^{1,2} topical application can also result in hypersensitivity.³

1. Chan CS, et al. Hydrocortisone-induced anaphylaxis. *Med J Aust* 1984; **141**: 444-6.2. Seale JP. Anaphylactoid reaction to hydrocortisone. *Med J Aust* 1984; **141**: 446.3. Corallo CE, Sosnin M. Bronchospasm, tachycardia following intravenous hydrocortisone. *Aust J Hosp Pharm* 1985; **15**: 103-4.4. Al Mahdy H, Hall M. Anaphylaxis and hydrocortisone. *Ann Intern Med* 1988; **108**: 487-8.5. Fulcher DA, Katalaris CH. Anaphylactoid reaction to intravenous hydrocortisone sodium succinate: a case report and literature review. *Med J Aust* 1991; **154**: 210-14.6. Kawane H. Anaphylactoid reaction to intravenous hydrocortisone sodium succinate. *Med J Aust* 1991; **154**: 782.7. Currie GP, et al. An unexpected response to intravenous hydrocortisone succinate in an asthmatic patient. *Br J Clin Pharmacol* 2005; **60**: 342.8. Wilkinson SM, et al. Hydrocortisone: an important cutaneous allergen. *Lancet* 1991; **337**: 761-2.

Interactions

The interactions of corticosteroids in general are described on p.1494.

Pharmacokinetics

For a brief account of the pharmacokinetics of corticosteroids, see p.1495.

Hydrocortisone is readily absorbed from the gastrointestinal tract and peak blood concentrations are attained in about an hour. The plasma half-life is about 100 minutes. It is more than 90% bound to plasma proteins. Following intramuscular injection, the absorption of the water-soluble sodium phosphate and sodium

succinate esters is rapid, while absorption of hydrocortisone free alcohol and its lipid-soluble esters is slower. Absorption of hydrocortisone acetate after intra-articular or soft-tissue injection is also slow. Hydrocortisone is absorbed through the skin, particularly in denuded areas.

Hydrocortisone is metabolised in the liver and most body tissues to hydrogenated and degraded forms such as tetrahydrocortisone and tetrahydrocortisol. These are excreted in the urine, mainly conjugated as glucuronides, with a very small proportion of unchanged hydrocortisone. Hydrocortisone readily crosses the placenta.

Uses and Administration

Hydrocortisone is a corticosteroid with both glucocorticoid and to a lesser extent mineralocorticoid activity (p.1490). As cortisol it is the most important of the predominantly glucocorticoid steroids secreted by the adrenal cortex. Hydrocortisone is used, usually with a more potent mineralocorticoid, for replacement therapy in adrenocortical insufficiency (p.1498). It may also be used for its glucocorticoid properties in other conditions for which corticosteroid therapy is indicated (p.1495) but drugs with fewer mineralocorticoid effects tend to be preferred for the long-term systemic therapy of auto-immune and inflammatory disease.

The dose may be expressed in terms of the base, and the following are each equivalent to about 100 mg of hydrocortisone:

- hydrocortisone acetate 112 mg
- hydrocortisone buteprate 135 mg
- hydrocortisone butyrate 119 mg
- hydrocortisone cipionate 134 mg
- hydrocortisone hydrogen succinate 128 mg
- hydrocortisone sodium phosphate 134 mg
- hydrocortisone sodium succinate 134 mg
- hydrocortisone valerate 123 mg

However, esterification generally alters potency and compounds with equivalent hydrocortisone content may not have equivalent clinical effect.

When given orally hydrocortisone free alcohol is usually used; the cipionate ester is used in some formulations. For replacement therapy in acute or chronic adrenocortical insufficiency the normal requirement is 20 to 30 mg daily (usually taken in 2 doses, the larger in the morning and the smaller in the early evening, to mimic the circadian rhythm of the body). Children may be given 400 to 800 micrograms/kg daily in 2 or 3 divided doses, adjusted as needed. Additional sodium chloride may be required if there is defective aldosterone secretion, but mineralocorticoid activity is usually supplemented by fludrocortisone acetate orally. Similar regimens have also been used to correct glucocorticoid deficiency in the salt-losing form of congenital adrenal hyperplasia (p.1502).

Hydrocortisone may be given intravenously, by slow injection or infusion, in the form of a water-soluble derivative such as hydrocortisone sodium succinate or hydrocortisone sodium phosphate when a rapid effect is required in emergencies: such conditions are acute adrenocortical insufficiency caused by Addisonian or post-adrenalectomy crises, by the abrupt accidental withdrawal of therapy in corticosteroid-treated patients, or by the inability of the adrenal glands to cope with increased stress in such patients; certain allergic emergencies such as anaphylaxis; acute severe asthma (status asthmaticus—see also p.1108); and shock. The usual dose is the equivalent of 100 to 500 mg of hydrocortisone, repeated 3 or 4 times in 24 hours, according to the severity of the condition and the patient's response. Children up to 1 year of age may be given 25 mg, those aged 1 to 5 years 50 mg, and those aged 6 to 12 years 100 mg. Fluids and electrolytes should be given as necessary to correct any associated metabolic disorder. Similar doses to those specified above may also be given intramuscularly but the response is like-

ly to be less rapid than that observed after intravenous doses. Corticosteroids are considered to be of secondary value in anaphylactic shock because of their relatively slow onset of action, but intravenous hydrocortisone may be a useful adjunct to adrenaline to prevent further deterioration in severely affected patients.

In patients with adrenal deficiency states supplementary corticosteroid therapy may be necessary during some surgical operations and hydrocortisone sodium succinate or sodium phosphate may be given intramuscularly or intravenously before surgery. Various regimens have been proposed (see also Surgery, p.1497). In patients taking more than 10 mg of oral prednisolone or its equivalent daily, the BNF recommends the following regimen:

- minor surgery under general anaesthesia, either the usual oral corticosteroid dose on the morning of surgery or hydrocortisone 25 to 50 mg (usually as the sodium succinate) intravenously at induction; the usual oral corticosteroid dose is resumed after surgery
- moderate or major surgery, the usual oral corticosteroid dose on the morning of surgery, plus hydrocortisone 25 to 50 mg intravenously at induction, and followed by similar doses of hydrocortisone 3 times daily, for 24 hours after moderate surgery and 48 to 72 hours after major surgery; the usual corticosteroid dose is resumed once hydrocortisone injections are stopped.

For local injection into soft tissues hydrocortisone is usually used in the form of the sodium phosphate or sodium succinate esters; doses in terms of hydrocortisone are usually 100 to 200 mg. For intra-articular injection hydrocortisone acetate is usually used in doses of 5 to 50 mg depending upon the size of the joint.

For topical application in the treatment of various skin disorders hydrocortisone and the acetate, buteprate, butyrate, and valerate esters are normally employed in creams, ointments, or lotions. Concentrations usually used have ranged from 0.1 to 2.5%. Although it is considered that hydrocortisone has fewer adverse effects on the skin and is less liable to cause adrenal suppression than the more potent topical corticosteroids (see p.1497 for a rough guide to the clinical potencies of topical corticosteroids), it should be borne in mind that this property may be considerably modified both by the type of formulation or vehicle used and by the type of esterification present; other factors that may also influence the degree of absorption include the site of application, use of an occlusive dressing, the degree of skin damage, and the size of the area to which the preparation is applied.

Hydrocortisone or its esters are also available in a variety of other dosage forms including those for ophthalmic, aural, dental, and rectal application, for use in allergic and inflammatory disorders.

Other esters of hydrocortisone that have occasionally been used include the aceponate, glycyrrhetinate, and propionate. Esters such as the aceponate may show modified topical activity.

Preparations

BP 2008: Gentamicin and Hydrocortisone Acetate Ear Drops; Hydrocortisone Acetate and Neomycin Ear Drops; Hydrocortisone Acetate and Neomycin Eye Ointment; Hydrocortisone Acetate Cream; Hydrocortisone Acetate Injection; Hydrocortisone Acetate Ointment; Hydrocortisone and Clioquinol Cream; Hydrocortisone Acetate and Neomycin Cream; Hydrocortisone Cream; Hydrocortisone Ointment; Hydrocortisone Oromucosal Tablets; Hydrocortisone Sodium Phosphate Injection; Hydrocortisone Sodium Succinate Injection; Miconazole and Hydrocortisone Acetate Cream; Miconazole and Hydrocortisone Cream; Miconazole and Hydrocortisone Ointment;

USP 31: Chloramphenicol and Hydrocortisone Acetate for Ophthalmic Suspension; Chloramphenicol, Polymyxin B Sulphate, and Hydrocortisone Acetate Ophthalmic Ointment; Clioquinol and Hydrocortisone Cream; Clioquinol and Hydrocortisone Ointment; Colistin and Neomycin Sulphates and Hydrocortisone Acetate Otic Suspension; Hydrocortisone Acetate Cream; Hydrocortisone Acetate Injectable Suspension; Hydrocortisone Acetate Lotion; Hydrocortisone Acetate Ointment; Hydrocortisone Acetate Ophthalmic Ointment; Hydrocortisone Acetate Ophthalmic Suspension; Hydrocortisone and Acetic Acid Otic Solution; Hydrocortisone Butyrate Cream; Hydrocortisone Cream; Hydrocortisone Gel; Hydrocortisone Injectable Suspension; Hydrocortisone Lotion; Hydrocortisone Ointment; Hydrocortisone Rectal Suspension; Hydrocortisone Sodium Phosphate Injection; Hydrocortisone Sodium Succinate for Injection;

Hydrocortisone Tablets; Hydrocortisone Valerate Cream; Hydrocortisone Valerate Ointment; Neomycin and Polymyxin B Sulfates and Hydrocortisone Acetate Cream; Neomycin and Polymyxin B Sulfates and Hydrocortisone Acetate Ophthalmic Suspension; Neomycin and Polymyxin B Sulfates and Hydrocortisone Ophthalmic Suspension; Neomycin and Polymyxin B Sulfates and Hydrocortisone Otic Solution; Neomycin and Polymyxin B Sulfates and Hydrocortisone Otic Suspension; Neomycin and Polymyxin B Sulfates; Bacitracin Zinc, and Hydrocortisone Acetate Ophthalmic Ointment; Neomycin and Polymyxin B Sulfates; Bacitracin Zinc, and Hydrocortisone Ointment; Neomycin and Polymyxin B Sulfates; Bacitracin Zinc, and Hydrocortisone Ophthalmic Ointment; Neomycin and Polymyxin B Sulfates; Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment; Neomycin and Polymyxin B Sulfates; Gramicidin, and Hydrocortisone Acetate Cream; Neomycin Sulfate and Hydrocortisone Acetate Cream; Neomycin Sulfate and Hydrocortisone Acetate Lotion; Neomycin Sulfate and Hydrocortisone Acetate Ointment; Neomycin Sulfate and Hydrocortisone Acetate Ophthalmic Ointment; Neomycin Sulfate and Hydrocortisone Acetate Ophthalmic Suspension; Neomycin Sulfate and Hydrocortisone Acetate Otic Suspension; Oxytetracycline Hydrochloride and Hydrocortisone Acetate Ophthalmic Suspension; Oxytetracycline Hydrochloride and Hydrocortisone Ointment; Polymyxin B Sulfate and Hydrocortisone Otic Solution.

Proprietary Preparations (details are given in Part 3)

Arg.: Alfacort; Anusol-HC; Azudridona; Cortenem†; Demacort; Eficort; Fridal; Hidrotiosa; Idenona†; Lactid HC; Locoid; Medrol; Microsan; Oralone; Schericur; Sirotimocin HC; Stiefcort; Transderma H; **Austral.:** Colifoam; Cortaid†; Cortef; Cortic; Derm-Aid; Egocort; Hycor; Hyson; Sigmacort; Siguient Hycor; Solu-Cortef; **Austria:** Colifoam; Eksenzialse F; Hydofatina sicc neumycin; Hydrocortone; Hydroderm; Locoidon; Retef; **Belg.:** Azacortine; Bucalsone†; Cortil†; Cremicort-H; Locoid; Nozema; Pannocort; Solu-Cortef; **Braz.:** Berilson; Cortisol; Cortizol; Cortizol; Cortizone; Hidrocort; Hidyn H; Locoid; Nutracort; Solu-Cortef†; Stefcort; Therasona; Westcor; **Canad.:** A-Hydrocort; Barriere-HC; Claritin Skin Itch Relief; Cortate; Cortef; Cortenema; Cortifoam; Cortoderm; Dermaflex HC; Dermalact Dicrict Anti-Itch; Emo-Cort; Hycort; Hyderm; Hydrodrene; HydroVal; Novo-Hydrocort; Prevex HC; Sarin HC; Solu-Cortef; Westcor; **Chile:** Aquanil H; Calmurd; Cortisol; Efficot; Hippo; Lactare-HC; Locoid; Nutracort; Nutralonat†; Pandel; Solu-Cortef; Vitulpast; **Cz.:** Laticort†; Locoid; Solu-Cortef†; **Denm.:** Colifoam; Locoid; Mildson; Solu-Cortef; Uniderm; **Fin.:** Ampiky; Apocort; Bucort; Colifoam; Kypakkas; Locoid; Nutracort†; Solu-Cortef; **Fr.:** Aphilan; Calmircort†; Colofam; Cortapsay; Cortisedermyl; Dermasemageon; Dermofenact†; Eficort; Hydracort; Locoid; MitoCortyl; Proctocort†; **Ger.:** Alfason; Colifoam; Dermallerg; Dermo Posteriort; Ebenol; Fenistil Hydrocort; Ficortril; Glycocortisol; Hydro-Wolf; Hydrocort; Hydrocutan; Hydrocutan-mild; Hydroderm HC; Hydrogenol; Laticort; Lindla Cort Hydro; Munirent; Pandel; Posterie Cortef; Posterie cort; ratioAllerg†; Remederm HC; Sanatson Mono; Soventol HC; Systral Hydrocort; velopural†; **Gr.:** Colifoam; Filocot; Lyo-Cort†; Nutracort†; Rolak; Solu-Cortef; **Hong Kong:** Cortef; Derm-Aid; Dhcort; Efficot†; Egocort; Hycort†; Hydrostone; Hytisone; Lacticare-HC; SigmaSmol; Solu-Cortef; **Hung.:** Laticort; Locoid; Solu-Cortef; **India:** Cipcorlin; Cutisoff; Entofoam; Wycort; **Indon.:** Berlicort; Calacort; Diakort; Lexacort; Loco; Steroderm; **Intl.:** Colifoam; Corian; Cortipon; Diadem; Hc45; Hydrocortisyl; Hydrocortone; Locoid; Mildson; Solu-Cortef; **Israel:** Cortifoam; Cortizone; Eficort; Lanacort; Solu-Cortef; **Ital.:** Colifoam; Colifoam; Cortil; Cortizol; Cortizone; Derm-Aid; Dhcort; Efficot; Egocort; Hycort†; Laticort; Locoid; Solu-Cortef; **Mex.:** Aquanil HC; Colicort; Eficort; Fadol; Flebocortid; Flemex; Icosan; **Neth.:** Buccalone; Cremicort; Locoid; Mildson; Solu-Cortef; **Norw.:** Colifoam; Locoid; Mildson; Solu-Cortef; **NZ:** BK HC; Colifoam; Derm-Aid; DP Hydrocortisone; Egocort; Lemnis Fatty Cream HC; Lipocort†; Locoid; Mildison; Skincalm; Solu-Cortef; **Philip.:** Cortiz; Droxiderm; Eficort; Hy-cort; Hydrotopic; Lacticare-HC; Pharmacort; Solu-Cortef; **Pol.:** Corhydron; Laticort; Locoid; Procriton; **Port.:** Colifoam; Dermimade Hidrocortisona†; Hidalone; Hydrocortone; Lactosana; Locoid; Pandel; Paraderm; Rapicort; Solu-Cortef; **Rus.:** Cortef (Кортеф); Laticort (Латикорт); Mylocort; Procutan; Solu-Cortef; Stopcort; **Singapore:** Derm-Aid; Dhcort; Efficot; Egocort; Hydrocort; Hydroderm; Solu-Cortef; Actocortina; Afasone; Ceneo; Dermosa Hidrocortisona; Hemodren; Hemorane; Hidroaltesona; Hidrocisidin; Isdnidrin; Lactisoma; Oral-sone; Scalpini; Capilar; Schericur; Suniderma; **Swed.:** Colifoam; Ficortri; Locoid; Mildson; Solu-Cortef; Uniderm; **Switz.:** Alfacortone; Glycosterone Ht; Hydrocortone; Locoid; Sandaniderm; Solu-Cortef; **Thail.:** Hytisone; Prevex HC; Solu-Cortef; **Turk.:** Cortimycine; Hipokort; Locoid; **UAE:** Alfa-cort; **UK:** Colifoam; Corlan; Cortipon; Corpertin; Demacort; Diadem; Efcoretan; Efcoresol; Exe-Cort; Hc45; Hydrocortisabt Hydrocortone; Lachacort; Locoid; Mildson; Solu-Cortef; Zenoxone; **USA:** A-Hydrocort; Aticort; Ala-Cort; Anuroc-HC; Aquanil HC; Bactine; CaldeCort; Carmol HC; Cetacort; Colocort; Cort-Dome; Cortaid; Cortef; Cortef Feminine Itch; Cortenema†; Corticaine; Cortifoam; Cortizone; Dermalact Dri-Cort; Dermol HC; Dermol HC; Esol-HC; Gynecort; Hemil-HC; Hi-Cor; Hydrocortone; HydroSkin; Hytone; Lacticare-HC; Lanacort; Locoid; Massengill Medicated; Neutrogena T/Scalp; Nutracort; Orbase HCA; Pandel; Penecort; Procor; Proctocort; Proctorem HC 25%; Recort Plus; Solu-Cortef; Synacort; Tegrin-HC; Texacort; U-Cort; Westcor; **Venez.:** Corticina Eff-cort; Hidrocort; Hidrozona; Histasona†; Lacticare-HC†; Liocort; Nutracort; Pantocort†; Solu-Cortef; Stricort.

Multi-ingredient: **Arg.:** Aeromicrona C†; Alercort†; Antibiocort; Anusol Duo; Anusol Duo S; Atodemura Plus; Bactisone; Bexon; Biocort; Cipro HC; Ciprocort; Ciproflox-Otic; Ciriax Otic; Ciriax Otic L; Colirio Antibiotico CNH; Cristalomicina; Delos Otic; Dercotex; Derivocort; Dermoperative; Disel Hidrocortisona; Epirhoc; Gentader; Gramicort†; Griseoprol; Hidrocortint; Hippoglos con Hidrocortisona; Ingal; Lactider HC†; Lidocort Proct; Linfol Dermico; Masivil Urea; Micozoj Compuesto†; Microsana C; Microsana Otic; Monizol Cort; Neu Pelvicin; Otex HC; Oto Biotaer; Otobiotic; Oticopiro; Otoperifit; Otoporin; Otoporin C; Otoporin L; Procto-Ikatral; Proctoret; Protoprocip; Prunisden Biotic†; Quemicetina con Hidrocortisona; Recaur†; Start NP†; Terra-Cortril; Terra-Cortril Nistatin†; Tocorectal; Tricur; Tridermal; Triefect†; Vagicurl; Xilader; Xyloprocto; **Austral.:** Ciproxin HC; Hydroform; Hydrozole; Proctosedyl; Rectinol HC; Resolv Plus; Xyloproct; **Austria:** C-Bildz Calmurd HC; Cortison Kemicetin; Hydofat; Hydrocortisone; Ichto-Cortin; Otoporin; Tropoderm; **Belg.:** Daktacort; Eoline†; Fucidin Hydrocortisone; Octose a l'Hydrocortisone; Otoporin†; Terra-Cortril; Terra-Cortril + Polymyxine B; **Braz.:** Anusol-HC; Cipro HC; Ginglone; Hemodott; Hidrocote; Hidroneo†; Oticinax; Otoporin; Terra-Cortril; Viiformio-Hidrocortisona; Xyloproct; **Canad.:** Anodan-HC; Anugesic-HC; Anusol-HC; Anuzinc HC; Anuzinc HC Plus; Cipro HC; Cortimyxin; Cortisporin; Egozinc-HC; Fucidin H; Neo-Cortef†; Pentamycetin-HC; PramoX HC; Proctodan-HC; Proctofoam-HC; Proctol; Proctomyxin HC; Proctosedyl; ratio-Hemicort-

HC; ratio-Proctosone; Rectogel HC; Rivasol HC; Sterex Plus; Uremol-HC; Vioform-Hydrocortisone; VoSol HC†; **Chile:** Cortifenol H†; Fucidin H; Otex HC; **Cz.:** Ciproxin HC Otic; Dobexil Plus†; Fucidin H; Ophthalmo-Framykin Compositum; Otoporin; Pimafofurt; Proctosedyl†; Septomixine; **Denm.:** Brentacort; Ciflox; Fucidin-Hydrocortison; Hydrocortison med Terramycin og Polymyxin-B; Locoidal; Proctosedyl; **Fin.:** Ciproxin-Hydrocortisone; Daktacon; Duocort; Fucidin-Hydrocortison; Locoidal†; Pantyon; Pimafofurt; Proctosedyl; Sibicort; Terra-Cortril; Terra-Cortril P; Trosycort; Xyloproct†; **Fr.:** Bacicoline; Madecassol Neomycine Hydrocortisone†; **Ger.:** Antiprurit†; Baycuten HC; Canesten HC†; Corti Jalkal†; Farco-Tril†; Fucidine plus†; Hydrodexan; Ichthocort; Novifort†; Nystadem comp; Piganorm; Polspectran HC; Posterisan Forte†; Terra-Cortril; **Gr.:** Ciprofloxacin HY†; Cortipheno H; Daktodor; Fucidin H; Hydrofusin; Terra-Cortril; **Hong Kong:** Anusol-HC†; Canesten HC; Cipro HC; Corticin†; Cortipheno H; Daktacort; Fucidin H; Hemcort HC†; Hydro-Funga; Micosone; Otoporin; Posterian Forte; Proctosedyl†; Proctosone†; Xyloproct; **Hung.:** Chlorocid-H†; Ciprobay†; Fucidin H; Otoporin; Oxycont; Pimafofurt; Posterian Forte; Tetran-Hydrocortison; **India:** Bell Diana Resolvent; Bell Resolvent; Belmycetin-C; Cortison Kemicetin†; Cortolam-m; Crotorax-HC; Daktacort; Eforlin; Furacin-S; Gentycin HC†; Keralin; Midithane; Multifungin H†; Neospiron-H; Pino-Cort; Proctosedyl; Shield; Wycont c Neomycin; **Indon.:** Anusol-HC; Binti; Chloramphecort; Dermacort; Epicorty; Haemocaine; Indoson; Kemiderm; Nufoart; Particol; Ramicort; Sanctorycin; Terra-Cortril; Thecort; Viydrohycort; Visancor; Zolacort; **Ir.:** Alphader; Alphysol HCF; Anugesic-HC; Anusol-HC; Calmurd HC; Canesten HC; Daktacort; Eurax-Hydrocortisone; Fucidin H; Gentisone HC; Locoid C; Nystafoma-HC†; Otoporin; Perinal; Proctofoa-HC; Proctosedyl; Quinocort†; Timodine; Vioform-Hydrocortisone; Xyloproct; **Israel:** Alphysol HC; Benzantine H; Calmurd HC†; Ciproxin HC; Daktacort; Epifo; Fucidin H; Hycocin; Hytisone; Hydrogastron; Panthisone; Perinal; Procto-Glyvenol; Proctofoam-HC; Proctorzon-N; Xyloproct†; **Ital.:** Argisone; Cort-Inal; Cortison Chemiceta; Emoril; Fucidin H; Idracemi; Idracemi Epiparia; Idrcotel†; Idroneomicil†; Kinogen; Mixtalone; Mobital; Nasomixin; Nevacort; Prepacor H; Preparazione Antiemorroridaria†; Protidol; Proctofoam-HC; Proctosedyl†; Proctosol; Reumacort; Scalcipin; Sedalen Cort†; Vasosterone; Vasosterone Antibiotico; Vasosterone Collirio; **Malaysia:** Beacort†; Canduc; Cipro HC†; Crotaniton H; Daktacort; Decortol; Foban-Hydro; Fucidin H; Miconazole H†; Neo-Hydro†; Pocin H; Proctosedyl†; Proctosone†; Setarin H†; Ucort; Xyloproct; Zaricort†; **Mex.:** Angenovag; Biotarant N; Ciproxina HC; Cloderm-H; Cortisporin; Daktacort; Dermanol; Hidrofenil; Hidroplicin; Litsei; Ofodox; Orel-Cl NF; Ofifar; Oto Eni; Otolone†; Poral; Soldrin; Sulfa Hidro; Ultracortin; Viiformo-Cort; Xyloderm; Xyloproct Plus; **Neth.:** Bacicoline-B; Calmurd HC; Daktacort; Otoporin; Pimafofurt; Proctosedyl; Terra-Cortril Gel Steracet met polymyxine-B†; Terra-Cortril met polymyxine-B†; **Norw.:** Daktacort; Fucidin-Hydrocortison; Locoidal; Proctosedyl; Terra-Cortril; Terra-Cortril Polymyxin B; Xyloproct; **NZ:** Ciproxin HC; Daktacort; DP Lotion H; Fungocort†; Locoid C; Micremc H; Pimafofurt; Proctosedyl; Xyloproct; **Philippines:** Cortsipor; Daktacort; Fucidin H; Hydrospor; Isonep H; Proctosedyl; Trimycin-H; **Pol.:** Atecorin; Chlornichalnid H; Daktacort†; Hemcort HC; Laticort-CH; Oxycont; Pimafofurt; Posterian H; Proctosone; **Port.:** Anuet; Clorcortic†; Cortil T; Daktacort; Davinicin†; Fuci-dine H; Leuco Hubber†; Locoid C†; Otoporin; Pimafofurt; Proctonostrum†; **Rus.:** Cortomyctin (Кортомицтин); Fucidin H (Фуцидин Г); Gioxysom (Гиоксион); Oxycont (Оксикорт); Pimafofurt (Пимафукорт); Posterian Forte (Постериан Фортэ); Relief Ultra (Реиф Ультра); **S.Afr.:** Calmurd HC†; Ciprobay HC; Daktacort; Fucidin H; Nasomixin†; Neoderm; Otoporin; Proctofoam†; Proctosedyl; Terra-Cortril; Viocont; **Singapore:** Candacon; Canesten HC; Ciprobay HC; Cortaid with Aloë†; Daktacort; Decortol; Fucidin H; Hydroderm-C; Micon-H; Neo-Hydro; Proctosedyl; Zaricort†; **Spain:** Afajuntav; Afasone B C; Anginovag; Anti-hemoroidal; Bacisporin; Brentan; Cilinofas Hidrocortisona; Ciproxina; Cohortan; Cortenema; Cortison Chemicet Topica; Dermo Hubber; Dentraine; Edifangine; Fuden H; Gritelat H; Gritelat Hidrocortisone†; Halibus Hidrocortisone; Hamodren Compuesto†; Hepru; Leuco Hubber; Milrosina Nistatina; Neo Hubber; Oto Difusor†; Oto Vitra†; Otoporin; Roberfarin; Terra-Cortril; Tisuderra; **Swed.:** Cortimycit (Кортимикит); Daktacort; Fenul-Hydrokortison; Fucidin-Hydrocortisone; Proctosedyl†; Terracortril; Terra-Cortril met polymyxine B; Xyloproct; Calmurd HC†; Ciproxin HC; Cortifuid N; Cortimycine; Daktacort; Demicalm-d; Fucidin H; Haemocort; Hydrocortisone compositum; Neo-Hydro; Otoporin; Septomixine; Terra-Cortril; **Thail.:** Antergan†; Candacon; Daktacort; Decortol; Dermal; Dorproct; Fucidin H; Ladocon; Otoporin; Proctosedyl; **Turk.:** Cormisin; Kortos; Ma-Ka-Ta; Ureacort; **UK:** Actinac; Alphader; Alphysol HC; Anugesic-HC; Anusol-HC; Plus HC; Calmurd HC; Canesten HC; Daktacort; Daktacort HC; Econacor; Eurax-Hydrocortisone; Fucidin H; Gentisone HC; Germoloids HC; Gregoderm†; Locoid C; Nystaform-HC; Otoporin; Perinal; Proctofoam-HC; Proctosedyl; Terra-Cortril†; Timodine; Uniroid-HC; Vioform-Hydrocortisone; Xyloproct; **USA:** I + 1-F; Acetasol HC; Alt-cort; Analpram-HC; AnaMantil HC; Anumed HC; Cipro HC; Coly-Mycin S Otic; Corgue; Cortane-B; Cortatigen; Cortic ND; Corticaine; Cortimycin; Cortisporin; Cortisporin-TC; Cortisporin†; Cytoc; Dermtex HC with Aloe; Ear-Eze; Emergent-Ez; Enzone; Epifoam; Fungold HC; HC Derma-Pax; HC Pramoxine; Hyson; LazerSporin-C; LidaMantle HC; Mediotic-HC; Neotrioxin HC; Novacort; Oticair†; Oti-Med†; Otic-Care; Oti-Tricin; Otobiotic; Otocort; Otomar-HC; Tomycin-HPN; Otoporin; Pediotic; Pramosone; Proctofoam-HC; Terra-Cortril; Tri-Otic†; UAD-Otic; Vanoxide-HC; VoSoL HC†; Vytone Xyralid; Zone-A; Zoto-HC; **Venez.:** Hipoglos con Hidrocortisona†; Ofalex; Quintic HC; Terra-Cortril†.


Isoflupredone Acetate (BANM, USAN, rINN) \otimes

Acetato de isoflupredona; 9 α -Fluoroprednisolone Acetate; Isoflupredone, Acetate d'; Isoflupredone Acetas; U-6013. 9 α -Fluoro-11 β ,17 α ,21-trihydroxyprogrena-1,4-diene-3,20-dione 21-acetate.

Изофлупредона Ацетат

$C_{23}H_{39}FO_6 = 420.5$

CAS — 338-95-4 (isoflupredone); 338-98-7 (isoflupredone acetate).



Pharmacopoeias. In US for veterinary use only.
USP 31 (Isoflupredone Acetate). Protect from light.

Profile

Isoflupredone acetate is a corticosteroid that has been used for its topical glucocorticoid activity (p.1490) in allergic rhinitis. Isoflupredone is also employed in veterinary medicine.

Preparations

Proprietary Preparations (details are given in Part 3)

Multi-ingredient: **Israel:** Proaf.

Loteprednol Etabonate (BANM, USAN, rINN) \otimes

CDDD-5604; Etabonato de loteprednol; HGP-1; Lotéprednol, Etabonate de; Loteprednol Ethyl Carbonate; Loteprednol Etabonas; P-5604. (11 β ,17 α -17-[Ethoxycarbonyl]oxy)-11-hydroxy-3-oxoandrosta-1,4-diene-17-carboxylic acid chloromethyl ester. Лотепреднола Этабонат

$C_{24}H_{31}ClO_7 = 467.0$.

CAS — 129260-79-3 (loteprednol); 82034-46-6 (loteprednol etabonate).

ATC — S01BA14.

ATC Vet — Q501BA14.

Profile

Loteprednol etabonate is a corticosteroid used for its glucocorticoid activity (p.1490) in the topical management of inflammatory and allergic disorders of the eye. It is usually used as eye drops containing 0.2 or 0.5%.

Prolonged application to the eye of preparations containing corticosteroids has caused raised intra-ocular pressure and reduced visual function.

◊ References.

1. Noble S, Goa KL. Loteprednol etabonate: clinical potential in the management of ocular inflammation. *BioDrugs* 1998; **10**: 329-39.

Preparations

Proprietary Preparations (details are given in Part 3)

Arg.: Alrex; Lored; Lotemax; Lotesoft; **Braz.:** Alrex†; **Ger.:** Lotemax; **Gr.:** Lotemax; **Hong Kong:** Lotemax; **India:** Loteflam; **Ital.:** Lotem; **Mex.:** Loterex; **Singapore:** Lotemax; **UK:** Lotemax; **USA:** Alrex; Lotemax; **Ven-
ez.:** Lotesoft.

Multi-ingredient: **Arg.:** Lotemicin.


Mazipredone (rINN) \otimes

Mazipredona; Maziprédone; Mazipredon. 11 β ,17-Dihydroxy-21-(4-methyl-1-piperazinyl)pregna-1,4-diene-3,20-dione.

Мазипредон

$C_{26}H_{38}N_2O_4 = 442.6$.

CAS — 13085-08-0.



Profile

Mazipredone is a corticosteroid used topically for its glucocorticoid activity (p.1490). It is used as the hydrochloride with miconazole in the treatment of fungal infections of the skin.

When applied topically, particularly to large areas, when the skin is broken, or under occlusive dressings, corticosteroids may be absorbed in sufficient amounts to cause systemic effects (p.1490). The effects of topical corticosteroids on the skin are